

---

## Plan Overview

*A Data Management Plan created using DMPonline*

**Title:** A cosmogenic-nuclide approach to quantifying the drivers of rocky coastline erosion under changing climate and sea-level conditions.

**Creator:** Abdoulie Faal

**Principal Investigator:** Dr Gordon Bromley

**Data Manager:** Gregor Maximilian Rink, Abdoulie Faal

**Project Administrator:** Dr Gordon Bromley

**Contributor:** Gregor Maximilian Rink

**Affiliation:** University College Dublin

**Template:** Health Research Board DMP Template

### Project abstract:

Large parts of the Irish coastline show rocky structures like cliffs and shore platforms (European Commission, 2023) and the effects of a rising sea level and erosion of shorelines and cliffs, including short term events like rockfalls and landslides, are potential geohazards for coastal areas (Violante, 2009; Yonggang et al., 2016). Despite large parts of the Ireland coastline are hard rock coasts (European Commission, 2023), a lot of the past research was regarding erosion and morphological changes of softer coasts, like beaches and estuaries (e.g. Carter and Stone, 1989; Cooper et al., 2004; Cronin et al., 2007; Swift et al., 2007; Guisado-Pintado and Jackson, 2019; Jackson et al., 2022) and much less research was focused on rocky coastlines (e.g. Thébaudeau et al., 2013; Cullen and Bourke, 2018; Cullen et al., 2018).

To fill this gap of knowledge, this research project will investigate erosion rates of rocky coasts by measuring the amount of produced  $^{10}\text{Be}$  in shore platforms, following an approach similar to Regard et al., 2012; Hurst et al., 2016, 2017 and Swirad et al., 2020. Due to a production rate of  $5.1 \pm 0.3$ , the measured amount  $^{10}\text{Be}$  reflects a long-term erosion rate in 1000 or 10000 year timescales (Gosse and Phillips, 2001; Markl, 2015). For studying the erosion history of the coastline on shorter timescales of years and decades and to investigate influences of storm events or seasons, this study uses historical map data, aerial photographs and Structure from Motion (SfM). SfM is a photogrammetric approach, that provides high resolution results in both temporal and spatial scales, while occurring as a fast and low-cost method with little gear requirements (Westoby et al., 2012; Gonçalves and Henriques, 2015; Piermattei et al., 2015; Swirad et al., 2019).

To gain as much information as possible, several sites along the Irish coast with different specifications in geology, geomorphology and geographical setting will be sampled and surveyed. Studying past coastal erosion and its drivers in different setting and locations will help us to model and simulate potential future evolutions of the Irish coastline, especially regarding climate change.

### References:

Carter, R.W.G., Stone, G.W., 1989. Mechanisms associated with the erosion of sand dune

cliffs, Magilligan, Northern Ireland. *Earth Surf. Process. Landforms* 14, 1–10. <https://doi.org/10.1002/esp.3290140102>

Cooper, J.A.G., Jackson, D.W.T., Navas, F., McKenna, J., Malvarez, G., 2004. Identifying storm impacts on an embayed, high-energy coastline: examples from western Ireland. *Marine Geology* 210, 261–280. <https://doi.org/10.1016/j.margeo.2004.05.012>

Cronin, K.M., Devoy, R.J.N., Gault, J., 2007. Modelling Estuarine Morphodynamics on the South Coast of Ireland. *Journal of Coastal Research* 474–479.

Cullen, N.D., Bourke, M.C., 2018. Clast abrasion of a rock shore platform on the Atlantic coast of Ireland: Clast abrasion of a rock shore platform. *Earth Surf. Process. Landforms* 43, 2627–2641. <https://doi.org/10.1002/esp.4421>

Cullen, N.D., Verma, A.K., Bourke, M.C., 2018. A comparison of structure from motion photogrammetry and the traversing micro-erosion meter for measuring erosion on shore platforms. *Earth Surface Dynamics* 6, 1023–1039. <https://doi.org/10.5194/esurf-6-1023-2018>

European Commission, 2023. EMODnet Map Viewer [WWW Document]. URL <https://emodnet.ec.europa.eu/geoviewer/#> (accessed 5.9.23).

Gonçalves, J.A., Henriques, R., 2015. UAV photogrammetry for topographic monitoring of coastal areas. *ISPRS Journal of Photogrammetry and Remote Sensing* 104, 101–111. <https://doi.org/10.1016/j.isprsjprs.2015.02.009>

Gosse, J.C., Phillips, F.M., 2001. Terrestrial in situ cosmogenic nuclides: theory and application. *Quaternary Science Reviews* 20, 1475–1560. [https://doi.org/10.1016/S0277-3791\(00\)00171-2](https://doi.org/10.1016/S0277-3791(00)00171-2)

Guisado-Pintado, E., Jackson, D.W.T., 2019. Coastal Impact From High-Energy Events and the Importance of Concurrent Forcing Parameters: The Cases of Storm Ophelia (2017) and Storm Hector (2018) in NW Ireland. *Front. Earth Sci.* 7, 190. <https://doi.org/10.3389/feart.2019.00190>

Hurst, M.D., Rood, D.H., Ellis, M.A., 2017. Controls on the distribution of cosmogenic  $^{10}\text{Be}$  across shore platforms. *Earth Surf. Dynam.* 5, 67–84. <https://doi.org/10.5194/esurf-5-67-2017>

Hurst, M.D., Rood, D.H., Ellis, M.A., Anderson, R.S., Dornbusch, U., 2016. Recent acceleration in coastal cliff retreat rates on the south coast of Great Britain. *Proc. Natl. Acad. Sci. U.S.A.* 113, 13336–13341. <https://doi.org/10.1073/pnas.1613044113>

Jackson, D.W.T., Short, A.D., Loureiro, C., Cooper, J.A.G., 2022. Beach morphodynamic classification using high-resolution nearshore bathymetry and process-based wave modelling. *Estuarine, Coastal and Shelf Science* 268, 107812. <https://doi.org/10.1016/j.ecss.2022.107812>

Markl, G., 2015. 4.8.3.9 Kosmogene Radionuklide, in: *Minerale und Gesteine: Mineralogie – Petrologie – Geochemie*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 565–569.

Piermattei, L., Carturan, L., Guarnieri, A., 2015. Use of terrestrial photogrammetry based on structure-from-motion for mass balance estimation of a small glacier in the Italian alps. *Earth Surface Processes and Landforms* 40, 1791–1802. <https://doi.org/10.1002/esp.3756>

Regard, V., Dewez, T., Bourlès, D.L., Anderson, R.S., Duperret, A., Costa, S., Leanni, L., Lasseur, E., Pedoja, K., Maillet, G.M., 2012. Late Holocene seacliff retreat recorded by  $^{10}\text{Be}$  profiles across a coastal platform: Theory and example from the English Channel. *Quaternary Geochronology* 11, 87–97. <https://doi.org/10.1016/j.quageo.2012.02.027>

Rink, G.M., Farrell, E.J., Bromley, G.R.M., 2024. Quantifying Aperiodic Cliff Top and Cliff Face Retreat Rates for an Eroding Drumlin on Ireland’s Atlantic Coast Using Structure-from-Motion. *Geosciences* 14, 165. <https://doi.org/10.3390/geosciences14060165>

Swift, L., Devoy, R., Wheeler, A., Sutton, G., Gault, J., 2007. Sedimentary dynamics and coastal changes on the south coast of Ireland. *J. Coast. Res.*

Swirad, Z.M., Rosser, N.J., Brain, M.J., 2019. Identifying mechanisms of shore platform erosion using Structure-from-Motion (SfM) photogrammetry. *Earth Surf. Process. Landforms* 44, 1542–1558. <https://doi.org/10.1002/esp.4591>

Swirad, Z.M., Rosser, N.J., Brain, M.J., Rood, D.H., Hurst, M.D., Wilcken, K.M., Barlow, J., 2020. Cosmogenic exposure dating reveals limited long-term variability in erosion of a rocky coastline. *Nat Commun* 11, 3804. <https://doi.org/10.1038/s41467-020-17611-9>

Thébaudeau, B., Trenhaile, A.S., Edwards, R.J., 2013. Modelling the development of rocky shoreline profiles along the northern coast of Ireland. *Geomorphology* 203, 66–78. <https://doi.org/10.1016/j.geomorph.2013.03.027>

Violante, C., 2009. Rocky coast: geological constraints for hazard assessment. *SP 322*, 1–31. <https://doi.org/10.1144/SP322.1>

Westoby, M.J., Brasington, J., Glasser, N.F., Hambrey, M.J., Reynolds, J.M., 2012. ‘Structure-from-Motion’ photogrammetry: A low-cost, effective tool for geoscience applications. *Geomorphology* 179, 300–314. <https://doi.org/10.1016/j.geomorph.2012.08.021>

Yonggang, J., Chaoqi, Z., Liping, L., Dong, W., 2016. Marine Geohazards: Review and Future Perspective. *Acta Geologica Sinica - English Edition* 90, 1455–1470. <https://doi.org/10.1111/1755-6724.12779>

**ID:** 164806

**Start date:** 01-09-2022

**End date:** 31-08-2026

**Last modified:** 23-12-2024

**Grant number / URL:** 13/RC/2092\_2

**Copyright information:**

The above plan creator(s) have agreed that others may use as much of the text of this plan as they would like in their own plans, and customise it as necessary. You do not need to credit the creator(s) as the source of the language used, but using any of the plan's text does not imply that the creator(s) endorse, or have any relationship to, your project or proposal

# A cosmogenic-nuclide approach to quantifying the drivers of rocky coastline erosion under changing climate and sea-level conditions.

## Data description and collection or re-use of existing data

### How will new data be collected or produced and/or how will existing data be re-used?

New data will consist of experimental data (lab measurements), field data (drone flights and GNSS) and outputs of drone analysis (point clouds and Digital Surface Models) and of numerical modelling. Existing data will consist of published experimental data from previous studies and field data, purchased from Tailte Éireann.

### What data (for example the kind, formats, and volumes), will be collected or produced?

Type of data	How will data be collected? (If re-using data indicate source)	Purpose of data collection?	File format(s)	Volume
experimental (new)	lab measurements	measuring concentrations of Beryllium-10	.txt	1-2 KB
experimental (existing)	reuse of existing Swirad et al. 2020, 10.1038/s41467-020-17611-9 Jeong et al. 2024 , 10.1016/j.margeo.2024.107291	setting up and testing the model of Swirad et al. (2020), before using it with own data	.txt	200KB
field data (new and output)	field measurements with drones and GNSS	generating high resolution Digital Surface Models for numerical modelling and coastal change analysis; generating 3D point clouds of costal cliffs for change analysis	.txt .png .e57 .tiff	500GB
field data (existing)	reuse of historical maps, purchased from Tailte Éireann.	Analysing of historical coastal change	wms-layers for GIS	
numerical (output)	numerical models	analysing possible coastal erosion based on measured concentrations of Beryllium-10	.mat .txt	1MB

## Documentation and data quality

### What metadata and documentation (for example the methodology of data collection and way of organising data) will accompany data?

Indicate which metadata will be provided to help others identify and discover the data.

Indicate which metadata standards (for example DDI, TEI, EML, MARC, CMDI) will be used and potential community standards available.

Use community metadata standards where these are in place

### What data quality control measures will be used?

Experimental data: standardised workflow, 1 blank for every five samples

field data: standardised workflow, calibration of measured coordinates via GNSS base station

numerical data: use of an already reviewed model (Swirad et al. 2020, 10.1038/s41467-020-17611-9)

## Storage and backup during the research process

### How will data and metadata be stored and backed up during the research process?

All data and metadata will be stored on the computer and automatically uploaded to the cloud system of the University of Galway and backed up on an external hard drive at least once every month. Research output will be backed up at the google-drive folder provided by iCrag.

### How will data security and protection of sensitive data be taken care of during the research?

Detail the key risks to the confidentiality and security related to human participants or other sensitive data and how this information will be managed.

## Legal and ethical requirements, codes of conduct

### If personal data are processed, how will compliance with legislation on personal data and on security be ensured?

No personal data will be processed

### How will other legal issues, such as intellectual property rights and ownership, be managed? What legislation is applicable?

Explain who will be the owner of the data and who will have the rights to control access

### What ethical issues and codes of conduct are there, and how will they be taken into account?

Consider whether ethical issues can affect how data are stored and transferred, who can see or use them, and how long they are kept.

## Data sharing and long-term preservation

### How and when will data be shared? Are there possible restrictions to data sharing or embargo reasons?

Explain how the data will be discoverable and shared (for example by deposit in a trustworthy data repository, indexed in a catalogue, use of a secure data service, direct handling of data requests, or use of another mechanism).

### How will data for preservation be selected, and where data will be preserved long-term (for example a data repository or archive)?

Indicate what data must be retained or destroyed for contractual, legal, or regulatory purposes.

### What methods or software tools are needed to access and use data?

Indicate whether potential users need specific tools to access and (re-)use the data. Consider the sustainability of software needed for accessing the data.

**How will the application of a unique and persistent identifier (such as a Digital Object Identifier (DOI)) to each data set be ensured?**

Explain how the data might be re-used in other contexts. Persistent identifiers should be applied so that data can be reliably and efficiently located and referred to. Persistent identifiers also help to track citations and re-use.

## **Data management responsibilities and resources**

**Who (for example role, position, and institution) will be responsible for data management (i.e. the data steward)?**

Outline the roles and responsibilities for data management/stewardship activities for example data capture, metadata production, data quality, storage and backup, data archiving, and data sharing. Alongside the PI, specify who is responsible for ensuring of the completion of these tasks.

**What resources (for example financial and time) will be dedicated to data management and ensuring that data will be FAIR (Findable, Accessible, Interoperable, Re-usable)?**

Explain how the necessary resources (for example time) to prepare the data for sharing/preservation (data curation) have been costed in. Carefully consider and justify any resources needed to deliver the data. These may include storage costs, hardware, staff time, costs of preparing data for deposit, and repository charges.